

Nutrition and Health Info-Sheet

For Health Professionals

Produced by
Karrie Heneman, PhD
Sheri Zidenberg-Cherr, PhD
UC Cooperative Extension
Center for Health and
Nutrition Research
Department of Nutrition
University of California
Davis, CA 95616-8669
October 2008

Some Facts About Flavonols

What are flavonols?

Flavonols are phytochemical compounds found in high concentrations in a variety of plant-based foods and beverages. Based on their structure, flavonols are classified as flavonoids and include the following compounds: quercetin, kaempferol, and myricetin. The specific amounts of flavonols in foods are affected by a range of factors including plant type and growth, season, light, degree of ripeness, food preparation, and processing. Despite these variables, high concentrations of flavonols can be found in apples, apricots, beans, broad beans, broccoli, cherry tomatoes, chives, cranberries, kale, leeks, pear, onions, red grapes, sweet cherries, and white currants (1).



What is the flavanol content of some common foods? (2)

<i>Food</i>	<i>Quercetin Content (mg/100g)</i>	<i>Kaempferol (mg/100g)</i>	<i>Myricetin (mg/100g)</i>
Apples	4.27	0.02	0.00
Apricots	2.08	0.00	0.00
Beans	0.00	31.32	0.00
Broad Beans (Fava Beans)	0.55	0.35	0.00
Broccoli	2.51	0.01	4.01
Cherry Tomatoes	2.76	0.10	0.00
Chives	4.77	10.00	0.00
Cranberries	15.09	0.09	6.78
Kale	7.71	26.74	0.00
Leeks	0.10	2.95	0.00
Pear	4.51	0.00	0.00
Onions	21.42	0.62	0.02
Red Grapes	1.38	0.00	0.01
Sweet Cherries	2.64	0.00	0.00
Tea	2.74	0.88	0.89
White Currant	2.68	0.17	0.18

Are there beneficial effects associated with consumption of flavonols?

Consumption of flavonols has been associated with a variety of beneficial effects including increased activity of erythrocyte superoxide dismutase (an antioxidant enzyme found in red blood cells), a decrease in lymphocyte DNA damage, a decrease in urinary 8-hydroxy-2'-deoxyguanosine (a marker of oxidative damage), and an increase in plasma antioxidant capacity (the ability to scavenge free radicals) (3).

There seems to be a lot of media hype around apples, broccoli, and cranberries. Are these really "super foods"?

As shown in the above table, apples, broccoli, and cranberries are all high in flavonols, in addition to a variety of other polyphenolic compounds. Below is a summary of the research that has looked at the effects of consuming these foods:

- **Apples:** Studies have investigated the relationship between consumption of apples and susceptibility to chronic diseases such as cancer, cardiovascular disease, asthma, and diabetes (4).
 - **Cancer:** A reduced risk of oral, pharynx, esophagus, colon, larynx, breast, ovary, and prostate cancer has been observed in individuals who consume greater than or equal to one apple a day in comparison to those who consume less (5). In support of these findings, previous research on *in vitro* systems has found that apple flavonoids can positively affect carcinogen bioactivation, cell-signaling, cell cycle regulation, angiogenesis (the formation of blood vessels), oxidative stress, and inflammation (5).
 - **Cardiovascular Disease:** Consumption of apples has been associated with a reduced risk of cardiovascular disease, cardiovascular events, coronary mortality, and thrombotic stroke (4).
 - **Asthma:** In adults, a reduced risk of asthma, bronchial sensitivity, and chronic obstructive pulmonary disease in addition to improved lung function (as measured by forced expiratory volume) has been associated with apple consumption (4).
 - **Diabetes:** Apple consumption has also been associated with a reduced risk of type 2 diabetes. In comparison to women who did not consume apples, consumption of at least one apple per day was associated with a 28% reduced risk of this chronic disease (6).
 - **Conclusions:** Current research supports an inverse relationship between consumption of apples and a variety of chronic diseases. In light of these findings, consuming an apple a day as part of your recommended intake of fruits and vegetables may prove to be beneficial to overall health; however, variety is important and apples should not be the sole fruit consumed by an individual.
- **Broccoli:** Cruciferous vegetables are part of the plant family Brassicaceae, which includes broccoli, in addition to cabbage, cauliflower, Brussels sprouts, turnips, and watercress. Consumption of this group of plant foods has been associated with a reduction in risk of several cancers including lung, breast, colorectal, and prostate (7).



- **Cancer:** Epidemiological evidence supports an inverse association between consumption of broccoli and breast cancer risk in pre-menopausal women (8), prostate cancer (9), bladder cancer in men (10), lung cancer in postmenopausal women, colon cancer in postmenopausal

women, cancer of the respiratory tract, stomach cancer in men, cancer of the reproductive organs in women, and thyroid cancer (11).

- *Conclusions:* In light of this research, the American Cancer Society recommends consuming broccoli (in any form, although raw broccoli contains the most nutrients) as part of a balanced diet that includes foods from a variety of plant sources.
- **Cranberries:** Cranberries are commonly touted as a remedy for treating urinary tract infections. Current research has also investigated the relationship between consumption of cranberry products and cancer and cardiovascular disease.
 - *Urinary Tract Infections:* Current scientific evidence suggests that use of cranberry products can prevent new urinary tract infections, possibly through inhibition of bacterial adhesion to the urinary tract (12). However, caution is warranted as cranberry juice may interact with warfarin (a blood thinner), resulting in bleeding (13).
 - *Cancer:* Despite promising findings in experimental models suggesting that cranberries can protect against breast, colon, prostate, lung, and other tumors (14), current human research does not provide evidence that consumption of cranberry products reduces risk of cancer (15).
 - *Heart Disease:* There is limited clinical evidence showing that consumption of cranberry products can protect against cardiovascular disease, possibly by decreasing LDL oxidation and increasing plasma antioxidant capacity (16).
 - *Conclusions:* The Dietary Guidelines for Americans, 2005 recommends limiting consumption of juice to ensure adequate intake of fiber (31g/ 2000 kcals). For women prone to urinary tract infections, consuming ½ cup of cranberry juice daily may help to reduce the number of new infections. As with apples and broccoli, current scientific evidence supports consumption of cranberry products as part of a varied diet.



References:

1. Aheme SA, O'Brien NM. Dietary flavonols: chemistry, food content, and metabolism. *Nutrition* 2002;18:75-81.
2. Nutrient Data Laboratory US Department of Agriculture. USDA Database for the Flavonoid Content of Selected Foods. US Department of Agriculture, Agricultural Research Service, 2007.
3. Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr* 2005;81:243S-255S.
4. Boyer J, Liu RH. Apple phytochemicals and their health benefits. *Nutr J* 2004;3:5.
5. Gallus S, Talamini R, Giacosa A, et al. Does an apple a day keep the oncologist away? *Ann Oncol* 2005;16:1841-4.

6. Song Y, Manson JE, Buring JE, Sesso HD, Liu S. Associations of dietary flavonoids with risk of type 2 diabetes, and markers of insulin resistance and systemic inflammation in women: a prospective study and cross-sectional analysis. *J Am Coll Nutr* 2005;24:376-84.
7. Juge N, Mithen RF, Traka M. Molecular basis for chemoprevention by sulforaphane: a comprehensive review. *Cell Mol Life Sci* 2007;64:1105-27.
8. Ambrosone CB, McCann SE, Freudenheim JL, Marshall JR, Zhang Y, Shields PG. Breast cancer risk in premenopausal women is inversely associated with consumption of broccoli, a source of isothiocyanates, but is not modified by GST genotype. *J Nutr* 2004;134:1134-8.
9. Kirsh VA, Peters U, Mayne ST, et al. Prospective study of fruit and vegetable intake and risk of prostate cancer. *J Natl Cancer Inst* 2007;99:1200-9.
10. Michaud DS, Spiegelman D, Clinton SK, Rimm EB, Willett WC, Giovannucci EL. Fruit and vegetable intake and incidence of bladder cancer in a male prospective cohort. *J Natl Cancer Inst* 1999;91:605-13.
11. Verhoeven DT, Goldbohm RA, van Poppel G, Verhagen H, van den Brandt PA. Epidemiological studies on brassica vegetables and cancer risk. *Cancer Epidemiol Biomarkers Prev* 1996;5:733-48.
12. Jepson RG, Craig JC. A systematic review of the evidence for cranberries and blueberries in UTI prevention. *Mol Nutr Food Res* 2007;51:738-45.
13. Pham DQ, Pham AQ. Interaction potential between cranberry juice and warfarin. *Am J Health Syst Pharm* 2007;64:490-4.
14. Neto CC. Cranberry and its phytochemicals: a review of in vitro anticancer studies. *J Nutr* 2007;137:186S-193S.
15. Duthie SJ, Jenkinson AM, Crozier A, et al. The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers. *Eur J Nutr* 2006;45:113-22.
16. Ruel G, Pomerleau S, Couture P, Lamarche B, Couillard C. Changes in plasma antioxidant capacity and oxidized low-density lipoprotein levels in men after short-term cranberry juice consumption. *Metabolism* 2005;54:856-61.

** Production of this material was supported by a grant from the Vitamin Cases Consumer Settlement Fund, created as a result of an antitrust class action. One of the purposes of the fund is to improve the health and nutrition of California consumers.*

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or service in the uniformed services (as defined by the Uniformed Services Employment and Reemployment Rights Act of 1994: service in the uniformed services includes membership, application for membership, performance of service, application for service, or obligation for service in the uniformed services) in any of its programs or activities.

University policy also prohibits reprisal or retaliation against any person in any of its programs or activities for making a complaint of discrimination or sexual harassment or for using or participating in the investigation or resolution process of any such complaint.

University policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Equal Opportunity Director, University of California, Agriculture and Natural Resources, 1111 Franklin Street, 6th Floor, Oakland, CA 94607, (510) 987-0096.